

NUCLEON-NUCLEON SCATTERING IN THE $1/N_c$ EXPANSION

SILAS R. BEANE

*Department of Physics, University of Washington,
Seattle, WA 98195*

E-mail: sbeane@phys.washington.edu

The nucleon-nucleon ${}^3S_1 - {}^3D_1$ coupled-channel problem is solved analytically to leading order in a joint expansion in the quark masses and in $1/N_c$. An approximate expression is derived for the ${}^3S_1$ scattering length in the large- N_c limit, and the large- N_c behavior of the deuteron is discussed.

1 Introduction

When the number of colors, N_c , of the QCD gauge group is taken large, QCD simplifies. One may then hope to learn about the real world with $N_c = 3$, in an expansion in $1/N_c$ ^{1,2}. The relevance of the large- N_c expansion for nuclear physics was first discussed in the seminal paper by Witten, where it was argued that the nucleon-nucleon (NN) potential grows as N_c in the large- N_c limit, and Hartree theory becomes exact². In a recent revival of interest, it was shown that large- N_c nuclear interactions are spin-flavor $SU(4)$ symmetric, with Wigner's phenomenologically successful supermultiplet symmetry following as an accidental symmetry³. Subsequently, the large- N_c scaling of the NN potential was analyzed in a general way and large- N_c expectations were shown to explain generic features of the NN interaction⁴. Given the levels of complexity that must be unravelled in going from QCD to a theory of nuclear forces, it is quite remarkable that large- N_c counting rules work so well. Hence large- N_c methods offer an important link between QCD and nuclear physics. The establishment of such links is, of course, an important goal for nuclear physics. In this paper, basic aspects of NN scattering in the large- N_c approximation will be reviewed and the ${}^3S_1 - {}^3D_1$ NN coupled-channel system will be analyzed in the large- N_c limit. I will also say some words about the deuteron and the bound-state spectrum in the large- N_c limit.

Assuming confinement, in the large- N_c limit the QCD chiral symmetry is spontaneously broken⁵. With two light flavors the chiral symmetry breaking pattern, $U(2) \otimes U(2) \rightarrow U(2)$, gives rise to four Goldstone bosons, the pions and η . While the chiral symmetry breaking scale, Λ_χ , and the meson masses scale as N_c^0 , the baryon masses scale as N_c . In the large- N_c limit the nucleons and Δ s are degenerate and fall into an infinite-dimensional representation of a contracted $SU(4)$ spin-flavor symmetry⁶. In this paper I will assume that η

and the Δ s can be integrated out of the low-energy effective field theory (EFT) of large- N_c QCD in a manner consistent with the large- N_c expansion⁷. This is, of course, a highly non-trivial assumption. Consider the resulting EFT involving nucleons and pions below Λ_χ (for reviews, see Ref.⁸). This EFT consists of nucleons emitting and absorbing pions with an interaction highly constrained by chiral symmetry. The most general Lagrange density consistent with QCD symmetries is

$$\begin{aligned} \mathcal{L} = & N^\dagger \left[i\partial_t + \vec{\nabla}^2/(2M) \right] N - \frac{1}{2} \left[C_0^S (N^\dagger N)^2 + C_0^T (N^\dagger \vec{\sigma} N)^2 \right] \\ & + \frac{F_\pi^2}{4} \text{Tr} \partial_\mu \Sigma^\dagger \partial^\mu \Sigma + \omega \frac{F_\pi^2}{2} \text{Tr} \mathcal{M}_q (\Sigma^\dagger + \Sigma) + g_A N^\dagger \vec{A} \cdot \vec{\sigma} N + \dots \end{aligned} \quad (1)$$

where $\vec{A}$ is the axial-vector current and the ellipses represent operators with more derivatives, more powers of the quark mass matrix, and more nucleon fields. In the large- N_c limit, $F_\pi \propto \sqrt{N_c}$, $g_A \propto N_c$ and $\omega \propto N_c^0$ ². The four-nucleon operators scale as $C_0^T \propto 1/N_c$ and $C_0^S \propto N_c$ ³. When considering the interactions of more than one nucleon, the ordering of the operators in Eq. (1) by way of a power-counting scheme is nontrivial. This is especially so in the ${}^3S_1 - {}^3D_1$ coupled-channel of NN scattering in an EFT with pions, which I will focus on in this paper. The presence of the deuteron, a near-threshold bound state, and the highly-singular tensor force imply complicated nonperturbative physics.

2 NN Scattering at Large- N_c

Defining a sensible large- N_c limit for NN scattering is nontrivial because the nucleon mass grows as N_c . There are several limits we may consider. The momentum-space Schrödinger equation is

$$\left(\frac{\hat{p}^2}{M} + \hat{V} \right) |\Psi\rangle = E |\Psi\rangle, \quad (2)$$

where M is the nucleon mass. At fixed velocity, the momentum transfer scales as N_c . Since the potential also scales as N_c , an overall factor of N_c scales out of the Hamiltonian and Hartree theory becomes exact². Evidently there is no EFT description in this limit as characteristic momenta quickly overcome Λ_χ and a description in terms of quarks and gluons becomes appropriate. At fixed momentum transfer, which is the limit relevant to EFT^a, the nucleon

^aIn principle, the momentum transfer can scale as an inverse power of N_c in the EFT, but then one is constrained to threshold.

kinetic energy is suppressed by $1/N_c$ while the potential energy grows as N_c . In this limit, the nucleons are always below threshold for scattering⁹. The static nucleons settle at the bottom of the potential well and nuclear matter forms a classical crystal⁴. The absence of a suitable large- N_c limit to describe scattering at fixed momentum transfer poses something of a dilemma. One way out is to focus on the large- N_c scaling properties of the NN potential and then to compare the resulting predictions with modern phenomenological NN potentials that fit phase shifts to high precision^{3,4}. This has proved highly successful. The question of whether the fixed momentum transfer limit exists has been called into question in interesting recent work^{10,11} which points out that there would appear to be violations of the counting rules for the potential arising from diagrams with multiple meson exchanges. This is an important unsolved puzzle.

3 The (Un)Coupled Channel Problem at Large- N_c

The most economical power-counting scheme for the $^3S_1$ – $^3D_1$ coupled channels is an expansion about the chiral limit¹². The pion-exchange part of the NN force in the chiral limit is purely tensor,

$$V_T(r) = -\frac{3\alpha_\pi}{r^3} + \mathcal{O}(m_\pi^2), \quad (3)$$

where

$$\alpha_\pi = \frac{g_A^2}{16\pi F_\pi^2}. \quad (4)$$

At leading order in the EFT expansion there is a local four-nucleon operator containing the effect of non-pionic short-distance physics which renormalizes the tensor force at short distances. The coefficient of this operator is $C_0 \equiv C_0^S + C_0^T = C_0^S + \mathcal{O}(1/N_c)$ ³. I choose to regulate this coordinate-space delta-function using a square well,

$$C_0 \delta^{(3)}(r) \rightarrow \frac{3C_0 \theta(R-r)}{4\pi R^3} \equiv V_0 \theta(R-r), \quad (5)$$

with strength V_0 and width R . The S -wave and D -wave components of the $J=1$ system, $u(r)$ and $w(r)$ respectively, are coupled by the tensor potential, V_T , defined in Eq. (3). The long-distance potential outside the square well is

$$\mathcal{V}_L(r) = \begin{pmatrix} 0 & -2\sqrt{2} MV_T(r) \\ -2\sqrt{2} MV_T(r) & 2MV_T(r) - 6/r^2 \end{pmatrix}, \quad (6)$$

while the short-distance potential inside the square well is

$$\mathcal{V}_S(r) = \begin{pmatrix} -MV_0 & 0 \\ 0 & -MV_0 - 6/r^2 \end{pmatrix} . \quad (7)$$

Defining Ψ to be

$$\Psi(r) = \begin{pmatrix} u(r) \\ w(r) \end{pmatrix} , \quad (8)$$

the regulated Schrödinger equation takes the compact form

$$\Psi''(r) + (k^2 + \mathcal{V}_L(r)\theta(r-R) + \mathcal{V}_S(r)\theta(R-r)) \Psi = 0. \quad (9)$$

One may ask whether there is a well-defined large- N_c limit. In Eq. (9), MV_T and MV_0 scale as N_c^2 , while the angular-momentum barrier scales as N_c^0 . In the EFT we treat the momentum transfer as fixed (N_c -independent). The total center-of-mass-energy is then suppressed by $1/N_c^2$ compared to the interaction energy. Here we will focus on this rather peculiar large- N_c limit. At leading order in the $1/N_c$ expansion, the coupled channel problem can be diagonalized via the similarity transformation $\Psi \rightarrow S^{-1}\Psi$ where

$$S = \begin{pmatrix} \sqrt{2} & -1/\sqrt{2} \\ 1 & 1 \end{pmatrix} . \quad (10)$$

Clearly ${}^3S_1 - {}^3D_1$ mixing is suppressed in the large- N_c limit. The general solution of Eq. (9) in the large- N_c limit is a linear combination of Bessel functions. Keeping the leading contributions I find

$$\begin{aligned} u(r) &= \sqrt{2} A r^{3/4} \cos \left(2\sqrt{\frac{6\alpha_\pi M}{r}} + \phi_0 \right) - \frac{B}{\sqrt{2}} r^{3/4} \cosh \left(2\sqrt{\frac{12\alpha_\pi M}{r}} + \delta_0 \right) \\ w(r) &= A r^{3/4} \cos \left(2\sqrt{\frac{6\alpha_\pi M}{r}} + \phi_0 \right) + B r^{3/4} \cosh \left(2\sqrt{\frac{12\alpha_\pi M}{r}} + \delta_0 \right), \end{aligned} \quad (11)$$

where ϕ_0 , δ_0 , A and B are determined from boundary conditions. Note that the repulsive hyperbolic solutions grow exponentially with N_c . Since these solutions cannot match to a power law in N_c , we find $B = 0$ in the large- N_c limit. We then have

$$u(r) = \sqrt{2} A r^{3/4} \cos \left(2\sqrt{\frac{6\alpha_\pi M}{r}} + \phi_0 \right) = \sqrt{2} w(r), \quad (12)$$

or

$$\frac{u(r)}{w(r)} = \sqrt{2} + O\left(\frac{1}{N_c^2}\right) \quad (13)$$

and the S -wave and D -wave wavefunctions oscillate *in phase*, with a ratio of $\sqrt{2}$. If there are nodes in the wavefunctions, then they must overlap. It is remarkable that such detailed information about nuclear wavefunctions can be deduced directly from QCD. Notice that the $1/N_c$ expansion is equivalent to the short-distance expansion. Eq. (13) is a remarkably accurate representation of the short-distance part of deuteron wavefunctions in conventional NN models with both hard- and soft-cores whose long-range part is governed by pion exchange¹³. At large distances in the real world, the angular momentum barrier is not negligible. With $M\alpha_\pi \equiv 1/\Lambda_{NN}$, the tensor force dominates the angular momentum barrier for distances $r \ll 1/\Lambda_{NN}$, which are outside the validity of the EFT description. It might prove useful to assign a more realistic large- N_c scaling to the angular momentum barrier and treat it as a perturbation using the WKB approximation, which becomes arbitrarily accurate in the large- N_c limit.

The phase ϕ_0 is determined by matching logarithmic derivatives of the interior and exterior wavefunctions,

$$\sqrt{-MV_0}R \cot\left(\sqrt{-MV_0}R\right) = \frac{3}{4} + \sqrt{\frac{6M\alpha_\pi}{R}} \tan\left(2\sqrt{\frac{6M\alpha_\pi}{R}} + \phi_0\right). \quad (14)$$

This equation determines the renormalization-group flow of the four-nucleon contact operator^{14,12}. The multi-branch structure of the right side of Eq. (14) is a consequence of the nonperturbative treatment of the tensor force. Treating $R \equiv 1/\Lambda$ as a Wilsonian cutoff, we see that the tensor force becomes perturbative when $\Lambda/\Lambda_{NN} \ll 1$. In the large- N_c limit, $\Lambda/\Lambda_{NN} \propto N_c^2 \gg 1$ and the tensor force becomes nonperturbative at arbitrarily small momentum scales. This suggests an interesting bound-state spectrum in the large- N_c limit.

4 A Deuteron in the Large- N_c Limit?

It is an intriguing mystery that QCD has a single scale, Λ_{QCD} , which is $\mathcal{O}(100 \text{ MeV})$, while characteristic nuclear binding energies are $\mathcal{O}(1 \text{ MeV})$. In particular, the deuteron binding energy, $-B$, is $B = 2.224575 \text{ MeV}$. The problem might not actually be this severe. The tensor force in the chiral limit is governed by a single scale, F_π . If we neglect geometrical factors and assume that the momenta of the bound nucleons is $\sim F_\pi$, then we would expect a

binding energy of order $F_\pi^2/M \sim 10$ MeV. Of course this argument breaks down as we vary N_c , since the axial coupling grows with N_c . Before considering the bound state spectrum in the pionful EFT, we will consider a simplified scenario.

At distance scales much larger than the pion Compton wavelength, the pion can be integrated out and the EFT simplifies considerably¹⁵. In EFT(π) observables are determined purely by four-nucleon contact operators⁸. In the ${}^3S_1 - {}^3D_1$ coupled-channel there is a single momentum-independent interaction, $C_0 \propto N_c$. We will assume that this operator is dominant at low energies in the large- N_c limit. The ${}^3S_1$ scattering length is then

$$a = R \left(1 - \frac{\tan(\sqrt{-MV_0}R)}{\sqrt{-MV_0}R} \right) \longrightarrow \frac{MC_0}{4\pi}, \quad (15)$$

where we use a square-well regulator and the arrow indicates that renormalization has been performed and C_0 is a renormalized quantity. Hence $a \propto N_c^2$. If the sign of C_0 is positive, then there is a bound state with binding energy

$$B = \frac{16\pi^2}{M^3 C_0^2}. \quad (16)$$

Hence, $B \propto 1/N_c^5$. What do we expect if we include pions in the EFT? We have seen that the tensor force dominates the potential at long distances. Hence we can write the Schrödinger equation in coordinate space as

$$\left(\frac{\nabla^2}{N_c \bar{M}} - \frac{3N_c \bar{\alpha}_\pi}{r^3} \right) |\Psi\rangle = E |\Psi\rangle, \quad (17)$$

where the N_c dependence has been scaled out and the barred quantities are N_c -independent. We can scale out the N_c dependence through a rescaling of the coordinate, $r \rightarrow N_c^2 \bar{r}$. The eigenvalue equation then becomes

$$\frac{1}{N_c^5} \left(\frac{\bar{\nabla}^2}{\bar{M}} - \frac{3\bar{\alpha}_\pi}{\bar{r}^3} \right) |\Psi_B\rangle = B |\Psi_B\rangle, \quad (18)$$

and we again find $B \propto 1/N_c^5$. This argument can be formulated in a more physical way as follows¹⁶. We can write the potential as

$$V_T \sim \frac{N_c}{r_B^3}, \quad (19)$$

where r_B can be loosely interpreted as the size of the bound state and I have assumed that all dimensional parameters are given by powers of Λ_{QCD} , which

is defined to be unity. Now assume that there is a bound state in the $^3S_1$ channel at large- N_c . The presence of a bound state requires a balance between the kinetic energy, $N_c v_B^2/2$, and the potential energy, and therefore we expect $V_T \sim N_c v_B^2$. Here v_B is the velocity of the bound state. As required for consistency, N_c drops out of this relation and we arrive at $v_B^2 \sim r_B^{-3}$. This is simply a statement of the virial theorem. Since the nucleon momentum is $N_c v_B$, it follows from the uncertainty principle that $v_B \sim (N_c r_B)^{-1}$. Again using the virial theorem, we find

$$r_B \propto N_c^2 \quad v_B \propto 1/N_c^3 \quad B \propto 1/N_c^5, \quad (20)$$

and we have a bound state that is large, slow, and shallow in the large- N_c limit. Hence our naive scaling expectations in the pionful EFT agree with those from EFT($\not{f}$). Unfortunately both of these arguments would appear to be too naive as they ultimately rely on the virial theorem constraint that the coordinate scales as N_c^2 , or, equivalently, that the momentum transfer scales as $1/N_c^2$. As we will see below, a more detailed study in the pionful theory suggests that there are an infinite number of bound states in the large- N_c limit, with the ground state binding energy increasing as a complicated function of N_c . This last point is not surprising, given the large- N_c wavefunctions found in the previous section. If one acts with the kinetic energy operator on the wavefunction of Eq. (12), one finds a complicated (harmonic) dependence on N_c .

5 The Bound-State Spectrum

One way of studying the bound-state spectrum is via the scattering length. Although a scattering length is strictly speaking not defined for a $1/r^3$ potential¹⁴, we can impose an infrared cutoff, $\mathcal{X}$, on V_T and match to a wavefunction at $r > \mathcal{X}$ of the form $r - a^{3S_1}$. This gives

$$a^{3S_1} = \frac{\mathcal{X} \left(-1 + 4\sqrt{6M\alpha_\pi/\mathcal{X}} \tan \left(2\sqrt{6M\alpha_\pi/\mathcal{X}} + \phi_0 \right) \right)}{\left(3 + 4\sqrt{6M\alpha_\pi/\mathcal{X}} \tan \left(2\sqrt{6M\alpha_\pi/\mathcal{X}} + \phi_0 \right) \right)}. \quad (21)$$

The purpose of this formula is to give analytical understanding of the bound state spectrum in the large- N_c limit. We would expect the infrared cutoff to take a value around $1/m_\pi$. Choosing an ultraviolet cutoff $R = 0.375$ fm and a square-well depth of $MV_0 = -2.07 \times 10^6$ MeV² which reproduce the scattering length for the physical value of $N_c = 3$ in the full problem, gives $\phi_0 = -7.6$. I then choose the infrared cutoff $\mathcal{X} = 0.97$ fm to reproduce the scattering length in the approximate formula. In order to explore the




Figure 1: The $^3S_1$ scattering length as a function of N_c . The solid line is a numerical solution of the Schrödinger equation with a small but finite pion mass and with $R = 0.375$ fm and a square-well depth of $MV_0 = -2.07 \times 10^6$ MeV² tuned to the physical scattering length at $N_c = 3$ ($\phi_0 = -7.6$). The dashed line is the approximate analytic solution, Eq. (21) with $\phi_0 = -7.6$ and $\mathcal{X} = 0.97$ fm which reproduce the physical scattering length at $N_c = 3$.

bound state spectrum as a function of N_c , I naively scale all quantities in the approximate formula and in the exact numerical result by the expected large- N_c scaling. The approximate formula for the scattering length and the exact numerical result are plotted as a function of N_c in Fig. (1). The approximate formula is reasonably accurate in the vicinity of $N_c = 3$ (for several branches of the tangent). The corresponding plot of the binding energy is given in Fig. (2). The plotted expression is derived from the approximate formula for the scattering length using effective range theory, and has been verified numerically along the plateaus using the exact solution. With the given choice of cutoff and square-well depth, there are no nodes in the wavefunction and the deuteron (at $N_c = 3$) is, by construction, the ground state. As N_c is increased from its physical value, the deuteron binding energy grows as a complicated function of N_c and quickly passes beyond Λ_χ and out of the EFT. There is then no bound state in the EFT. As N_c approaches 4, a new bound state, the first excited state, appears at threshold and then undergoes the same decoupling as N_c is increased. This cycle repeats itself an infinite number of times in the large- N_c limit. The excited levels can be studied in more detail through the singularities of the scattering length, using the approximate formula, Eq. (21). One can estimate the density of eigenstates¹⁷ for the $1/r^3$ potential using WKB; with excitation number, n , one finds $|E_n| \sim n^6 \Lambda_{QCD}$ with $n = 1, 2, \dots$. Therefore as $N_c \rightarrow \infty$ there are an infinite number of bound states, infinitely




Figure 2: The deuteron binding energy as a function of N_c abstracted from the approximate analytical solution, Eq. (21), and verified numerically along the plateaus.

dispersed in energy. But there would appear never to be more than one bound state present in the EFT.

6 Caveats

There are many caveats to the large- N_c picture of NN scattering presented in this paper. Among them is the absence of the Δ degrees of freedom. NN scattering becomes a significantly more complicated coupled channel problem when the Δ s are degenerate with the nucleons⁷. It would be interesting to repeat the analysis presented here with the complete contracted spin-flavor $SU(4)$ multiplet. A more serious problem lies in the fact that the large- N_c limit in which the momentum transfer is held fixed does not experience scattering, and potentially does not exist as a sensible limit of QCD^{10,11}. One may also seriously question the procedure of varying N_c slightly away from its physical value to deduce information about the large- N_c limit. It is conceivable that the true large- N_c Schrödinger equation behaves quite differently. On a related note, there are several issues of noncommutativity of limits. First there is the issue of whether to take the large- N_c limit at the level of the Schrödinger equation, or at the level of its solution. There is also the issue of whether the chiral limit commutes with the large- N_c limit; note that the tensor force in the chiral limit is highly singular at all distances, and this has an important effect on the bound state spectrum. Anthropic arguments suggest that the fine-tuned nature of the deuteron binding energy is essential to the emergence of life. The results presented in this paper suggest that the large- N_c limit can

be relevant to nature only at discrete values of N_c where there continues to be a bound state with deuteron quantum numbers and binding energy within anthropic bounds. Whether this be the ground state or the N_c th excited state is irrelevant in the EFT. Other (large) values of N_c –away from the plateaus in Fig. (2)– would in all likelihood not support the intelligent life necessary to think about large- N_c QCD.

Acknowledgments

I am grateful for valuable conversations with David Kaplan, Markus Luty, Martin Savage and Bira van Kolck. Special thanks to the organizers, particularly Rich Lebed, for a stimulating workshop. This research was supported in part by the DOE grant DE-FG03-97ER41014.

References

1. G. 't Hooft, *Nucl. Phys. B* **72**, 461 (1974).
2. E. Witten, *Nucl. Phys. B* **160**, 57 (1979).
3. D.B. Kaplan and M.J. Savage, *Phys. Lett. B* **365**, 244 (1996).
4. D.B. Kaplan and A.V. Manohar, *Phys. Rev. C* **56**, 76 (1997).
5. S. Coleman and E. Witten, *Phys. Rev. Lett.* **45**, 100 (1980).
6. R.F. Dashen and A.V. Manohar, *Phys. Lett. B* **315**, 425 (1993).
7. M.J. Savage, *Phys. Rev. C* **55**, 2185 (1997).
8. U. van Kolck, *Prog. Part. Nucl. Phys.* **43**, 337 (1999); S.R. Beane, P.F. Bedaque, W.C. Haxton, D.R. Phillips and M.J. Savage, Essay for the Festschrift in honor of Boris Ioffe, in the 'Encyclopedia of Analytic QCD', edited by M. Shifman, by World Scientific; [hep-ph/0008064](#).
9. S. Coleman, *Aspects of Symmetry*, Cambridge University Press, 1985.
10. M.K. Banerjee, T.D. Cohen and B.A. Gelman, *Phys. Rev. C* **65**, 034011 (2002).
11. A.V. Belitsky and T.D. Cohen, [hep-ph/0202153](#).
12. S.R. Beane, P.F. Bedaque, M.J. Savage and U. van Kolck, *Nucl. Phys. A* **700**, 377 (2002).
13. D.W.L. Sprung, W. van Dijk, E. Wang, D.C. Zheng, P. Sarriguren and J. Martorell, *Phys. Rev. C* **49**, 2942 (1994).
14. S.R. Beane, P.F. Bedaque, L. Childress, A. Kryjevski, J. McGuire and U. van Kolck, *Phys. Rev. A* **64**, 042103 (2001).
15. S. Weinberg, *Nucl. Phys. B* **363**, 3 (1991).
16. M.A. Luty, *private communication*.
17. A.M. Perelomov and V.S. Popov, *Teor. i. Mate. Fiz.* **4**, 48 (1970).